

EXHIBIT A

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

Product	Cat. No.	Size	Quantity	Price
20X SSC	15557-044	1 L	—	\$ 22.00
	15557-035	4 L	—	66.00
	15557-028	10 L	—	155.00

20X SSC (Ultra Pure) is a solution formulated for use in nucleic acid hybridizations and blot transfer applications. It is used in concentrations ranging from 0.2X to 20X, depending on the application. Supplied in 1-L plastic bottles, or in a 4-L or 10-L stackable CUBITAINER® Box, 20X SSC contains 3.0 M NaCl and 0.3 M sodium citrate, at pH 7.0.

Analytical Specifications

pH of 20X SSC at 23°C	7.0 ± 0.1
specific conductance of 20X SSC at 23°C	170 ± 10 mS/cm

Performance and quality testing: No detectable contaminating activity is observed in DNA nicking and ribonuclease assays.

Recommended storage condition: 15°C to 30°C.

CUBITAINER® is a registered trademark of Hewlett Corporation, Inc.

20X SSPE	15591-043	1 L	—	\$ 27.00
	15591-035	4 L	—	81.00
	15591-027	10 L	—	189.00

20X SSPE (Ultra Pure) is a prepared buffer concentrate formulated for use in nucleic acid hybridizations and blot transfer applications. Supplied in 1-L plastic bottles, or in a 4-L or 10-L stackable CUBITAINER® Box, 20X SSPE contains 3.0 M NaCl, 0.2 M NaH₂PO₄, and 0.02 M EDTA, at pH 7.4.

Analytical Specifications

pH of 20X SSPE at 25°C	7.4 ± 0.1
specific conductance of 20X SSPE at 25°C	190 ± 10 mS/cm

Performance and quality testing: No detectable contaminating activity is observed in DNA nicking and ribonuclease assays.

Recommended storage condition: 15°C to 30°C.

CUBITAINER® is a registered trademark of Hewlett Corporation, Inc.

Streptavidin Agarose	15942-014	5 ml	—	\$195.00
-----------------------------	-----------	------	---	----------

For additional information about this product, please refer to page 14-9.

Sucrose	15503-014	1 kg	—	\$25.00
	15503-022	5 kg	—	69.00

Sucrose (Ultra Pure) is suitable for forming density gradients for a variety of separation applications. It is used to separate RNAs, to prepare bacteriophage λ DNA arms, and to separate proteins. Sucrose gradients typically are formed by use of gradient formers or by layering two or more sucrose solutions in varying amounts in a centrifuge tube.

Analytical Specifications

molecular weight	342.30
appearance	white, free-flowing crystals or powder
purity	≥99.9%
insolubles in a 50% (w/v) solution	none detected
lead	≤8 ppm
free glucose	≤0.1%

Performance and quality testing: No detectable contaminating activity is observed in protease assays.

Recommended storage condition: 15°C to 30°C, dry.

See also:

Gradient Formers, page 28-15.




EXHIBIT B

1. 10/1/2010
2. 10/1/2010
3. 10/1/2010
4. 10/1/2010
5. 10/1/2010
6. 10/1/2010
7. 10/1/2010
8. 10/1/2010
9. 10/1/2010
10. 10/1/2010
11. 10/1/2010
12. 10/1/2010
13. 10/1/2010
14. 10/1/2010
15. 10/1/2010
16. 10/1/2010
17. 10/1/2010
18. 10/1/2010
19. 10/1/2010
20. 10/1/2010
21. 10/1/2010
22. 10/1/2010
23. 10/1/2010
24. 10/1/2010
25. 10/1/2010
26. 10/1/2010
27. 10/1/2010
28. 10/1/2010
29. 10/1/2010
30. 10/1/2010
31. 10/1/2010
32. 10/1/2010
33. 10/1/2010
34. 10/1/2010
35. 10/1/2010
36. 10/1/2010
37. 10/1/2010
38. 10/1/2010
39. 10/1/2010
40. 10/1/2010
41. 10/1/2010
42. 10/1/2010
43. 10/1/2010
44. 10/1/2010
45. 10/1/2010
46. 10/1/2010
47. 10/1/2010
48. 10/1/2010
49. 10/1/2010
50. 10/1/2010
51. 10/1/2010
52. 10/1/2010
53. 10/1/2010
54. 10/1/2010
55. 10/1/2010
56. 10/1/2010
57. 10/1/2010
58. 10/1/2010
59. 10/1/2010
60. 10/1/2010
61. 10/1/2010
62. 10/1/2010
63. 10/1/2010
64. 10/1/2010
65. 10/1/2010
66. 10/1/2010
67. 10/1/2010
68. 10/1/2010
69. 10/1/2010
70. 10/1/2010
71. 10/1/2010
72. 10/1/2010
73. 10/1/2010
74. 10/1/2010
75. 10/1/2010
76. 10/1/2010
77. 10/1/2010
78. 10/1/2010
79. 10/1/2010
80. 10/1/2010
81. 10/1/2010
82. 10/1/2010
83. 10/1/2010
84. 10/1/2010
85. 10/1/2010
86. 10/1/2010
87. 10/1/2010
88. 10/1/2010
89. 10/1/2010
90. 10/1/2010
91. 10/1/2010
92. 10/1/2010
93. 10/1/2010
94. 10/1/2010
95. 10/1/2010
96. 10/1/2010
97. 10/1/2010
98. 10/1/2010
99. 10/1/2010
100. 10/1/2010

Harsh treatment: Pour several hundred milliliters of boiling 0.1% SDS onto the membrane. Cool to room temperature.

If a membrane is to be reprobed, it must not be allowed to dry out between hybridization and stripping. If it becomes dry, the probe may bind to the matrix.

2. Place membrane on a sheet of dry Whatman 3MM filter paper and blot excess liquid with a second sheet. Wrap the membrane in plastic wrap and set up an autoradiograph.

If signal is still seen after autoradiography, rewash using harsher conditions.

3. The membrane can now be rehybridized. Alternatively, it can be dried and stored for later use.

Membranes can be stored dry between Whatman 3MM paper for several months at room temperature. For long-term storage, place the membranes in a desiccator at room temperature or 4°C.

REAGENTS AND SOLUTIONS

Aqueous prehybridization/hybridization (APH) solution

5× SSC (APPENDIX 2)

5× Denhardt solution (APPENDIX 2)

1% (w/v) SDS

Add 100 µg/ml denatured salmon sperm DNA (see below) just before use

Alternatives to Denhardt solution and denatured salmon sperm DNA as blocking agents are listed in Table 2.10.5 (see discussion in critical parameters).

Denatured salmon sperm DNA

Dissolve 10 mg Sigma type III salmon sperm DNA (sodium salt) in 1 ml water. Pass vigorously through a 17-G needle 20 times to shear the DNA. Place in a boiling water bath for 10 min, then chill. Use immediately or store at -20°C in small aliquots. If stored, reheat to 100°C for 5 min and chill on ice immediately before using.

Formamide prehybridization/hybridization (FPH) solution

5× SSC (APPENDIX 2)

5× Denhardt solution (APPENDIX 2)

50% (w/v) formamide

1% (w/v) SDS

Add 100 µg/ml denatured salmon sperm DNA (see above) just before use

Alternatives to Denhardt solution and denatured salmon sperm DNA as blocking agents are listed in Table 2.10.5 (see discussion in critical parameters).

Commercial formamide is usually satisfactory for use. If the liquid has a yellow color, deionize as follows: add 5 g of mixed-bed ion-exchange resin [e.g., Bio-Rad AG 501-X8 or 501-X8(D) resins] per 100 ml formamide, stir at room temperature for 1 hr, and filter through Whatman no. 1 paper.

CAUTION: Formamide is a teratogen. Handle with care.

Labeling buffer

200 mM Tris-Cl, pH 7.5

30 mM MgCl₂

10 mM spermidine

Mild stripping solution

5 mM Tris-Cl, pH 8.0

2 mM EDTA

0.1× Denhardt solution (APPENDIX 2)

CURRENT PROTOCOLS IN MOLECULAR BIOLOGY

VOLUME 1

EDITORIAL BOARD

Frederick M. Ausubel

Massachusetts General Hospital & Harvard Medical School

Roger Brent

Massachusetts General Hospital & Harvard Medical School

Robert E. Kingston

Massachusetts General Hospital & Harvard Medical School

David D. Moore

Massachusetts General Hospital & Harvard Medical School

J.G. Seidman

Harvard Medical School

John A. Smith

University of Alabama

Kevin Struhl

Harvard Medical School

GUEST EDITORS

Lisa M. Albright

DNA Sequencing

Donald M. Coen

Harvard Medical School

Polymerase Chain Reaction

Ajit Varki

University of California San Diego

Glycoproteins

SERIES EDITOR

Virginia Benson Chanda



John Wiley & Sons, Inc.

CORE 13 (\$36)

Copyright © 1994–1997 by John Wiley & Sons, Inc.

Copyright © 1987–1994 by Current Protocols

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Section 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc.

While the authors, editors, and publisher believe that the specification and usage of reagents, equipment, and devices, as set forth in this book, are in accord with current recommendations and practice at the time of publication, they accept no legal responsibility for any errors or omissions, and make no warranty, express or implied, with respect to material contained herein. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of experimental reagents, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each chemical, piece of equipment, reagent, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. This is particularly important in regard to new or infrequently employed chemicals or experimental reagents.

Library of Congress Cataloging in Publication Data:

Current protocols in molecular biology. 3 vols.

1. Molecular biology—Technique. 2. Molecular biology—Laboratory manuals. I. Ausubel, Frederick M.


QH506.C87 1987 574.8'8'028 87-21033

ISBN 0-471-50338-X

Printed in the United States of America

20 19 18 17 16 15 14 13

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

improper page starting from 54

☐ *Scanned copy is best available.*

SCANNED # 20